



Recommended Adult* Equine Preventative Vaccination Program (1/16)
 (*Horses 1 year of Age or Older)



VACCINE	ADMINISTERED <i>IM - Intramuscular</i> <i>IN - Intranasal</i>	FREQUENCY
Tetanus/Sleeping Sickness ¹	IM	Annually ⁴
Tetanus/Sleeping Sickness/West Nile Virus Combination	IM	Annually ^{3&4}
Rabies	IM	Annually ⁴
Flu/Rhino or Flu	IM or IN	Every 6 months
Strangles (IN) ²	IN	Annually ⁴
West Nile	IM	Annually ^{4&5}

- ¹ In case of a penetrating wound, Tetanus Toxoid should be given if greater than 6 months since vaccination. If no prior Tetanus Toxoid can be verified, consider using Tetanus Antitoxin.
- ² We recommend the use of this vaccine only where there is known or possible exposure.
- ³ If this vaccine is given in late March to a previously vaccinated horse, a West Nile booster in the same year should not be necessary
- ⁴ Previously unvaccinated adults should be boosted 3-4 weeks after their initial vaccine.
- ⁵ This vaccine should be given in March or April (once yearly) if the Tetanus/Sleeping Sickness/West Nile combination is not used.

FOR MORE INFORMATION:

(916) 652 7645

CONNECT WITH US:

-  facebook.com/LBEMC
-  [@loomis_basin_equine](https://instagram.com/@loomis_basin_equine)

Limited time only. Some restrictions may apply. Call for complete details.